arXiv:hep-th/9405040v1 6 May 1994

MULTIMOMEMTUM HAMILTONIAN FORMALISM IN FIELD THEORY. GEOMETRIC SUPPLEMENTARY

Gennadi Sardanashvily

Department of Theoretical Physics, Moscow State University
117234 Moscow, Russia
E-mail: sard@theor.phys.msu.su

The well-known geometric approach to field theory is based on description of classical fields as sections of fibred manifolds, e.g. bundles with a structure group in gauge theory. In this approach, Lagrangian and Hamiltonian formalisms including the multiomentum Hamiltonian formalism are phrased in terms of jet manifolds [1, 2, 3, 5, 6, 8, 9, 10]. Then, configuration and phase spaces of fields are finite-dimensional. Though the jet manifolds have been widely used for theory of differential operators, the calculus of variations and differential geometry, this powerful mathematical methods remains almost unknown for physicists. This Supplementary to our previous article [10] aims to summarize necessary requisites on jet manifolds and general connections [7, 9, 11].

All morphisms throughout are differentiable mappings of class C^∞. Manifolds are real, Hausdorff, finite-dimensional, second-countable and connected.

We use the conventional symbols $\otimes$, $\vee$ and $\wedge$ for the tensor, symmetric and exterior products respectively. The interior product (contraction) is denoted by $\rfloor$.

The symbols ∂^A_B mean the partial derivatives with respect to coordinates with indices ${}^B_A$.

Given a manifold M with an atlas of local coordinates (z^λ), the tangent bundle TM of M (resp. the cotangent bundle T^*M of M) is provided with the atlas of the induced coordinates $(z^\lambda, \dot z^\lambda)$ (resp. $(z^\lambda, \dot z_\lambda)$ relative to the holonomic bases ∂_λ (resp. dz^λ).

If $f : M \to M'$ is a manifold mapping, by

$$Tf : TM \to TM' \qquad \dot z'^\lambda = \frac{\partial f^\lambda}{\partial z^\alpha}\dot z^\alpha,$$

is meant the morphism tangent to f.

By pr_1 and pr_2, we denote the canonical surjections

$$\mathrm{pr}_1 : A \times B \to A, \qquad \mathrm{pr}_2 : A \times B \to B.$$

We handle the following types of manifold mappings $f : M \to M'$ when the tangent morphism Tf to f meets the maximal rank. These are immersion, submersion and local diffeomorphism when f is both immersion and submersion.

Recall that a mapping $f : M \to M'$ is called immersion (resp. submersion) at a point $z \in M$ when the tangent morphism Tf to f is injection (resp. surjection) of the tangent space T_zM to M at z to the tangent space $T_{f(z)}M'$. A manifold mapping f of M is termed immersion (resp. submersion) if it is immersion (resp. submersion) at all points of M.

A triple $f : M \to M'$ is called the submanifold (resp. the fibred manifold) if f is both immersion and injection (resp. both submersion and surjection) of M to M'. A submanifold which also is a topological subspace is called the imbedded submanifold. Every open subset U of a manifold M is endowed with the manifold structure such that the canonical injection $i_U : U \hookrightarrow M$ is imbedding.

1 Fibred manifolds

Throughout the work, by Y is meant a fibred manifold

$$\pi : Y \to X \tag{1}$$

over an n-dimensional base X. We use symbols y and x for points of Y and X respectively.

The total space Y of a fibred manifold $Y \to X$, by definition, is provided with an atlas of fibred coordinates

$$\begin{aligned} &(x^\lambda, y^i), \qquad x^\lambda \circ \pi = x^\lambda, \\ &x^\lambda \to x'^\lambda(x^\mu), \qquad y^i \to y'^i(x^\mu, y^j), \end{aligned} \tag{2}$$

where $(x^\lambda$ are coordinates of X. They are compatible with the fibration (1).

A fibred manifold $Y \to X$ is called the locally trivial fibred manifold if there exists a fibred coordinate atlas of Y over an open covering $\{\pi^{-1}(U_\xi)\}$ of Y where $\{U_\xi\}$ is an open covering of the base X. In other words, all points of a fibre of Y belong to the same coordinate chart. By a differentiable fibre bundle (or simply a bundle), we mean the locally trivial fibred manifold (1) provided with a family of equivalent bundle atlases

$$\Psi = \{U_\xi, \psi_\xi\}, \qquad \psi_\xi : \pi^{-1}(U_\xi) \to U_\xi \times V,$$

where V is a standard fibre of Y. Recall that two bundle atlases are called equivalent if their union also is a bundle atlas.

If $Y \to X$ is a bundle, the fibred coordinates (2) of Y are assumed to be bundle coordinates associated with a bundle atlas Ψ of Y, that is,

$$y^i(y) = (\phi^i \circ \mathrm{pr}_2 \circ \psi_\xi)(y), \qquad \pi(y) \in U_\xi, \tag{3}$$

where ϕ^i are coordinates of the standard fibre V of Y.

Given fibred manifolds $Y \to X$ and $Y' \to X'$, by a fibred morphism is meant a fibre-to-fibre manifold mapping $\Phi : Y \to Y'$ over a manifold mapping $f : X \to X'$. If $f = \mathrm{Id}\,_X$, fibred morphism is termed the fibred morphism $\underset{X}{\to}$ over X.

In particular, let X_X denotes the fibred manifold $\mathrm{Id}\,_X : X \hookrightarrow X$. Given a fibred manifold $Y \to X$, a fibred morphism $X_X \to Y$ over X is a global section of $Y \to X$. It is a closed imbedded submanifold. Let N be an imbedded submanifold of X. A fibred morphism $N_N \to Y$ over $N \hookrightarrow X$ is called a section of $Y \to X$ over N. For each point $x \in X$, a fibred manifold, by definition, has a section over an open neighborhood of x.

Remark. In accordance with the well-known theorem, if a fibred manifold $Y \to X$ has a global section, every section of Y over a closed imbedded submanifold N of X is extended to a global section of Y due to the properties which are required of a manifold.

If fibred morphism $Y \to Y'$ over X is a submanifold, $Y \to X$ is called the fibred submanifold of $Y' \to X$. Fibred imbedding and fibred diffeomorphism are usually callled the monomorphism and the isomorphism respectively.

Given a fibred manifold $Y \to X$, every manifold mapping $f : X' \to X$ yields the pullback $f^*Y \to X'$ comprising the pairs

$$\{(y, x') \in Y \times X' \mid \quad \pi(y) = f(x')\}$$

together with the surjection $(y, x') \to x'$. Every section s of the fibred manifold $Y \to X$ defines the corresponding pullback section

$$(f^*s)(x') = ((s \circ f)(x'), x'), \qquad x' \in X',$$

of the fibred manifold $f^*Y \to X'$.

In particular, if a mapping f is a submanifold, the pullback f^*Y is called the restriction $Y\mid_{f(X')}$ of the fibred manifold Y to the submanifold $f(X') \subset X$.

The product of fibred manifolds $\pi : Y \to X$ and $\pi' : Y' \to X$ over X, by definition, is the total space of pullbacks

$$\pi^*Y' = \pi'^*Y = Y \underset{X}{\times} Y'.$$

A composite fibred manifold (or simply a composite manifold) is defined to be composition of surjective submersions

$$\pi_{\Sigma X} \circ \pi_{Y\Sigma} : Y \to \Sigma \to X. \tag{4}$$

It is the fibred manifold $Y \to X$ provided with the particular class of coordinate atlases:

$$(x^\lambda, \sigma^m, y^i) \tag{5}$$

$$x'^\lambda = f^\lambda(x^\mu), \qquad \sigma'^m = f^m(x^\mu, \sigma^n), \qquad y'^i = f^i(x^\mu, \sigma^n, y^j),$$

where (x^μ, σ^m) are fibred coordinates of the fibred manifold $\Sigma \to X$.

In particular, let $TY \to Y$ be the tangent bundle of a fibred manifold $Y \to X$. We have the composite manifold

$$TY \overset{T\pi}{\to} TX \to X. \tag{6}$$

Given the fibred coordinates (2) of Y, the corresponding induced coordinates of TY are $(x^\lambda, y^i, \dot{x}^\lambda, \dot{y}^i)$.

The tangent bundle $TY \to Y$ of a fibred manifold Y has the subbundle

$$VY = \mathrm{Ker}\, T\pi$$

which is called the vertical tangent bundle of Y. This subbundle is provided with the induced coordinates $(x^\lambda, y^i, \dot y^i)$.

The vertical cotangent bundle $V^*Y \to Y$ of Y, by definition, is the vector bundle dual to the vertical tangent bundle $VY \to Y$, it is not a subbundle of T^*Y.

With VY and V^*Y, we have the following exact sequences of bundles over a fibred manifold $Y \to X$:

$$0 \to VY \hookrightarrow TY \underset{Y}{\to} Y \underset{X}{\times} TX \to 0, \tag{7a}$$

$$0 \to Y \underset{X}{\times} T^*X \hookrightarrow T^*Y \to V^*Y \to 0. \tag{7b}$$

For the sake of simplicity, we shall denote the products

$$Y \underset{X}{\times} TX, \qquad Y \underset{X}{\times} T^*X$$

by the symbols TX and T^*X respectively. Different splittings

$$Y \underset{X}{\times} TX \underset{Y}{\to} TY, \qquad V^*Y \to T^*Y$$

of the exact sequences (7a) and (7b), by definition, correspond to different connections on a fibred manifold Y.

At the same time, there is the canonical bundle monomorphism

$$\overset{n}{\wedge} T^*X \underset{Y}{\otimes} V^*Y \underset{Y}{\hookrightarrow} \overset{n+1}{\wedge} T^*Y. \tag{8}$$

Let $\Phi : Y \to Y'$ be a fibred morphism over f. The tangent morphism $T\Phi$ to Φ reads

$$(\dot x'^\lambda, \dot y'^i) \circ T\Phi = (\partial_\mu f^\lambda \dot x^\mu, \partial_\mu \Phi^i \dot x^\mu + \partial_j \Phi^i \dot y^j). \tag{9}$$

It is both a linear bundle morphism over Φ and a fibred morphism over the tangent morphism Tf to f. Its restriction to the vertical tangent subbundle VY yields the vertical tangent morphism

$$V\Phi : VY \to VY',$$

$$\dot y'^i \circ V\Phi = \partial_j \Phi^i \dot y^j. \tag{10}$$

Vertical tangent bundles of fibred manifolds utilized in field theory meet almost always the following simple structure.

One says that a fibred manifold $Y \to X$ has vertical splitting if there exists the linear isomorphism

$$\alpha : VY \underset{Y}{\to} Y \underset{X}{\times} \overline{Y} \tag{11}$$

where $\overline{Y} \to X$ is a vector bundle. The fibred coordinates (2) of Y are called adapted to the vertical splitting (11) if the induced coordinates of the vertical tangent bundle VY take the form

$$(x^\mu, y^i, \dot y^i = \overline{y}^i \circ \alpha)$$

where $(x^\mu, y^i, \overline{y}^i)$ are bundle coordinates of $\overline{Y}$. In this case, transition functions $\dot y^i \to \dot y'^i$ between induced coordinate charts are independent on the coordinates y^i.

In particular, a vector bundle $Y \to X$ has the canonical vertical splitting

$$VY = Y \underset{X}{\times} Y. \tag{12}$$

An affine bundle Y modelled on a vector bundle $\overline{Y}$ has the canonical vertical splitting

$$VY = Y \underset{X}{\times} \overline{Y}. \tag{13}$$

Moreover, linear bundle coordinates of a vector bundle and affine bundle coordinates of an affine bundle are always adapted to these canonical vertical splittings.

We shall refer to the following fact.

Lemma 1.1: Let Y and Y' be fibred manifolds over X and $\Phi : Y \to Y'$ a fibred morphism over X. Let $V\Phi$ be the vertical tangent morphism to Φ. If Y' admits vertical splitting $VY' = Y' \times \overline{Y}$, then there exists the linear bundle morphism

$$\overline{V}\Phi : VY \underset{Y}{\to} Y \times \overline{Y} \tag{14}$$

over Y given by the coordinate expression

$$\overline{y}'^i \circ \overline{V}\Phi = \partial_j \Phi^i \dot y^j.$$

By differential forms (or symply forms) on a fibred manifold, we shall mean exterior, tangent-valued and pullback-valued forms.

Recall that a tangent-valued r-form on a manifold M is defined to be a section

$$\phi = \phi^\mu_{\lambda_1 \dots \lambda_r} dz^{\lambda_1} \wedge \dots \wedge dz^{\lambda_r} \otimes \partial_\mu$$

of the bundle $\overset{r}{\wedge} T^*M \otimes TM$. In particular, tangent-valued 0-forms are vector fields on M.

Remark. There is the 1:1 correspondence between the tangent-valued 1-forms on M and the linear bundle morphisms $TM \to TM$ or $T^*M \to T^*M$ over M

$$\begin{aligned} &\theta : M \to T^*M \otimes TM, \\ &\theta : T_zM \ni t \mapsto t\rfloor\theta(z) \in T_zM, \\ &\theta : T^*_zM \ni t^* \mapsto \theta(z)\rfloor t^* \in T^*_zM. \end{aligned} \tag{15}$$

For instance, Id_{TM} corresponds to the canonical tangent-valued 1-form on M:

$$\theta_M = dz^\lambda \otimes \partial_\lambda, \qquad \partial_\lambda \rfloor \theta_M = \partial_\lambda.$$

Let $\overset{r}{\wedge}\mathcal{T}^*(M)$ be the sheaf of exterior r-forms on M and $\mathcal{T}(M)$ the sheaf of vector fields on M. Tangent-valued r-forms on a manifold M constitute the sheaf $\overset{r}{\wedge}\mathcal{T}^*(M)\otimes\mathcal{T}(M)$. It is brought into the sheaf of graded Lie algebras with respect to the Frölicher-Nijenhuis (F-N) bracket.

The F-N bracket is defined to be the sheaf morphism

$$\overset{r}{\wedge}\mathcal{T}^*(M)\otimes\mathcal{T}(M)\times\overset{s}{\wedge}\mathcal{T}^*(M)\otimes\mathcal{T}(M)\to\overset{r+s}{\wedge}\mathcal{T}^*(M)\otimes\mathcal{T}(M),$$

$$\begin{aligned}[][\phi,\sigma] &= -(-1)^{|\phi||\sigma|}[\sigma,\phi]=[\alpha\otimes u,\beta\otimes v]\\ &= \alpha\wedge\beta\otimes[u,v]+\alpha\wedge\mathbf{L}_u\beta\otimes v-(-1)^{rs}\beta\wedge\mathbf{L}_v\alpha\otimes u\\ &\quad +(-1)^r(v\rfloor\alpha)\wedge d\beta\otimes u-(-1)^{rs+s}(u\rfloor\beta)\wedge d\alpha\otimes v,\\ &\quad \alpha\in\overset{r}{\wedge}\mathcal{T}^*(M),\quad \beta\in\overset{s}{\wedge}\mathcal{T}^*(M),\quad u,v\in\mathcal{T}(M),\end{aligned}$$

where $\mathbf{L}_u$ and $\mathbf{L}_v$ are the Lie derivatives of exterior forms. We have its coordinate expression

$$\begin{aligned}[][\phi,\sigma] &= (\phi^\nu_{\lambda_1\ldots\lambda_r}\partial_\nu\sigma^\mu_{\lambda_{r+1}\ldots\lambda_{r+s}}\\ &\quad -(-1)^{rs}\sigma^\nu_{\lambda_1\ldots\lambda_s}\partial_\nu\phi^\mu_{\lambda_{s+1}\ldots\lambda_{r+s}}-r\phi^\mu_{\lambda_1\ldots\lambda_{r-1}\nu}\partial_{\lambda_r}\sigma^\nu_{\lambda_{r+1}\ldots\lambda_{r+s}}\\ &\quad +(-1)^{rs}s\sigma^\mu_{\lambda_1\ldots\lambda_{s-1}\nu}\partial_{\lambda_s}\phi^\nu_{\lambda_{s+1}\ldots\lambda_{r+s}})dz^{\lambda_1}\wedge\ldots\wedge dz^{\lambda_{r+s}}\otimes\partial_\mu.\end{aligned}$$

Given a tangent-valued form ϕ, the Nijenhuis differential is defined to be the sheaf morphism

$$d_\phi:\sigma\mapsto d_\phi\sigma=[\phi,\sigma]. \tag{16}$$

In particular, if $\theta=u$ is a vector field, we have the Lie derivative of tangent-valued forms

$$\begin{aligned}\mathbf{L}_u\sigma=[u,\sigma] &= (u^\nu\partial_\nu\sigma^\mu_{\lambda_1\ldots\lambda_s}-\sigma^\nu_{\lambda_1\ldots\lambda_s}\partial_\nu u^\mu\\ &\quad +s\sigma^\mu_{\lambda_1\ldots\lambda_{s-1}\nu}\partial_{\lambda_s}u^\nu)dx^{\lambda_1}\wedge\ldots\wedge dx^{\lambda_s}\otimes\partial_\mu.\end{aligned}$$

The Nijehuis differential (16) can be extended to exterior forms σ by the rule

$$\begin{aligned}d_\phi\sigma=\phi\rfloor d\sigma+(-1)^r d(\phi\rfloor\sigma) &= (\phi^\nu_{\lambda_1\ldots\lambda_r}\partial_\nu\sigma_{\lambda_{r+1}\ldots\lambda_{r+s}}\\ &\quad +(-1)^{rs}s\sigma_{\lambda_1\ldots\lambda_{s-1}\nu}\partial_{\lambda_s}\phi^\nu_{\lambda_{s+1}\ldots\lambda_{r+s}})dz^{\lambda_1}\wedge\ldots\wedge dz^{\lambda_{r+s}}.\end{aligned} \tag{17}$$

In particular, if $\phi=\theta_M$, the familiar exterior differential

$$d_{\theta_M}\sigma=d\sigma$$

is reproduced.

On a fibred manifold $Y\to X$, we consider the following particular subsheafs of exterior and tangent-valued forms:

(i) exterior horizontal forms $\phi : Y \to \overset{r}{\wedge} T^*X$;

(ii) tangent-valued horizontal forms

$$\phi : Y \to \overset{r}{\wedge} T^*X \underset{Y}{\otimes} TY,$$
$$\phi = dx^{\lambda_1} \wedge \ldots \wedge dx^{\lambda_r} \otimes (\phi^\mu_{\lambda_1\ldots\lambda_r}\partial_\mu + \phi^i_{\lambda_1\ldots\lambda_r}\partial_i);$$

(iii) tangent-valued projectable horizontal forms

$$\phi = dx^{\lambda_1} \wedge \ldots \wedge dx^{\lambda_r} \otimes (\phi^\mu_{\lambda_1\ldots\lambda_r}(x)\partial_\mu + \phi^i_{\lambda_1\ldots\lambda_r}(y)\partial_i);$$

(iv) vertical-valued horizontal forms

$$\phi : Y \to \overset{r}{\wedge} T^*X \underset{Y}{\otimes} VY,$$
$$\phi = \phi^i_{\lambda_1\ldots\lambda_r} dx^{\lambda_1} \wedge \ldots \wedge dx^{\lambda_r} \otimes \partial_i.$$

Vertical-valued horizontal 1-forms on $Y \to X$ are termed the soldering forms.

By pullback-valued forms on a fibred manifold $Y \to X$, we mean the morphisms

$$Y \to \overset{r}{\wedge} T^*Y \underset{Y}{\otimes} TX, \tag{18}$$

$$Y \to \overset{r}{\wedge} T^*Y \underset{Y}{\otimes} T^*X. \tag{19}$$

Let us emphasize that the forms (18) are not tangent-valued forms and the forms (19) are not exterior forms on Y. In particular, we shall refer to the pullback $\pi^*\theta_X$ of the canonical form θ_X on the base X by π onto Y. This is a pullback-valued horizontal 1-form on $Y \to X$ which we denote by the same symbol

$$\theta_X : Y \to T^*X \underset{Y}{\otimes} TX, \qquad \theta_X = dx^\lambda \otimes \partial_\lambda. \tag{20}$$

Horizontal n-forms on a fibred manifold $Y \to X$ are called horizontal densities. We denote

$$\omega = dx^1 \wedge \ldots \wedge dx^n.$$

2 Jet manifolds

This Section briefs some notions of the higher order jet formalism and illustrates them by the ones of first and second order jet machineries.

We use the multi-index Λ, $\mid \Lambda \mid = k$ for symmetrized collections $(\lambda_1 ... \lambda_k)$. By $\Lambda + \lambda$ is meant the symmetrized collection $(\lambda_1 ... \lambda_k \lambda)$.

Definition 1.2: The k-order jet space J^kY of a fibred manifold $Y\to X$ is defined to comprise all equivalence classes j^k_xs, $x\in X$, of sections s of Y so that sections s and s' belong to the same class j^k_xs if and only if

$$\partial_\Lambda s^i(x)=\partial_\Lambda s'^i(x), \qquad 0\leq\mid\Lambda\mid\leq k.$$

In other words, sections of $Y\to X$ are identified by the first $k+1$ terms of their Teylor series at points of X.

Given fibred coordinates (2) of Y, the k-order jet space J^kY is provided with atlases of the adapted coordinates

$$(x^\lambda, y^i_\Lambda), \qquad 0\leq\mid\Lambda\mid\leq k.$$

They bring J^kY into a finite-dimensional smooth manifold satisfying the conditions which we require of a manifold. It possesses the composite fibration

$$J^kY\to J^{k-1}Y\to ...\to Y\to X.$$

In particular, the first order jet manifold (or simply the jet manifold) J^1Y of Y consists of the equivalence classes j^1_xs, $x\in X$, of sections s of Y so that different sections s and s' belong to the same class j^1_xs if and only if

$$Ts\mid_{T_xX}=Ts'\mid_{T_xX}.$$

In other words, sections $s\in j^1_xs$ are identified by their values $s^i(x)=s'^i(x)$ and values of their partial derivatives $\partial_\mu s^i(x)=\partial_\mu s'^i(x)$ at the point x of X.

There are the natural surjections

$$\pi_1: J^1Y\ni j^1_xs\mapsto x\in X, \tag{21}$$

$$\pi_{01}: J^1Y\ni j^1_xs\mapsto s(x)\in Y. \tag{22}$$

We have the composite manifold

$$\pi_1=\pi\circ\pi_{01}: J^1Y\to Y\to X.$$

The surjection (21) is a fibred manifold. The surjection (22) is a bundle. If $Y\to X$ is a bundle, so is the surjection (21).

The first order jet manifold J^1Y of Y is provided with the adapted coordinate atlases

$$(x^\lambda, y^i, y^i_\lambda), \tag{23}$$

$$(x^\lambda, y^i, y^i_\lambda)(j^1_xs)=(x^\lambda, s^i(x), \partial_\lambda s^i(x)),$$

$$y'^i_\lambda=(\frac{\partial y'^i}{\partial y^j}y^j_\mu+\frac{\partial y'^i}{\partial x^\mu})\frac{\partial x^\mu}{\partial x'^\lambda}. \tag{24}$$

A glance at the transformation law (24) shows that the bundle $J^1Y \to Y$ is an affine bundle. We call it the jet bundle. It is modelled on the vector bundle

$$T^*X \underset{Y}{\otimes} VY \to Y. \tag{25}$$

The second order jet manifold J^2Y of a fibred manifold $Y \to X$ is defined to be the union of all equivalence classes $j^2_x s$ of sections s of $Y \to X$ such that sections $s \in j^2_x s$ are identified by their values and values of their first and second order partial derivatives at the point $x \in X$. The second order jet manifold J^2Y is endowed with the adapted coordinates

$$(x^\lambda, y^i, y^i_\lambda, y^i_{\lambda\mu} = y^i_{\mu\lambda}),$$

$$y^i_\lambda(j^2_x s) = \partial_\lambda s^i(x), \qquad y^i_{\lambda\mu}(j^2_x s) = \partial_\mu\partial_\lambda s^i(x).$$

Let Y and Y' be fibred manifolds over SXS and $\Phi : Y \to Y'$ a fibred morphism over a diffeomorphism f of X. It yields the k-order jet prolongation

$$J^k\Phi : J^kY \ni j^k_x s \mapsto j^k_{f(x)}(\Phi \circ s \circ f^{-1}) \in J^kY'$$

of Φ. In particular, the first order jet prolongation (or simply the jet prolongation) of Φ reads

$$\begin{gathered} J^1\Phi : J^1Y \to J^1Y', \\ J^1\Phi : j^1_x s \mapsto j^1_{f(x)}(\Phi \circ s \circ f^{-1}), \\ y'^i_\lambda \circ J^1\Phi = \partial_\lambda(\Phi^i \circ f^{-1}) + \partial_j(\Phi^i y^j_\lambda \circ f^{-1}). \end{gathered} \tag{26}$$

It is both an affine bundle morphism over Φ and a fibred morphism over the diffeomorphism f.

Every section s of a fibred manifold $Y \to X$ admits the k-order jet prolongation to the section

$$(J^ks)(x) \overset{\text{def}}{=} j^k_x s$$

of the fibred manifold $J^kY \to X$. In particular, its first order jet prolongation to the section J^1s of the fibred jet manifold $J^1Y \to X$ reads

$$(x^\lambda, y^i, y^i_\lambda) \circ J^1s = (x^\lambda, s^i(x), \partial_\lambda s^i(x)). \tag{27}$$

Every vector field

$$u = u^\lambda\partial_\lambda + u^i\partial_i$$

on a fibred manifold $Y \to X$ has the jet lift to the projectable vector field

$$\begin{aligned} &\overline{u} = r \circ J^1u : J^1Y \to TJ^1Y, \\ &\overline{u} = u^\lambda\partial_\lambda + u^i\partial_i + (\partial_\lambda u^i + y^j_\lambda\partial_j u^i - y^i_\mu\partial_\lambda u^\mu)\partial^\lambda_i, \end{aligned} \tag{28}$$

on the fibred jet manifold $J^1Y \to X$. To construct it, we use the canonical fibred morphism

$$r : J^1TY \to TJ^1Y,$$

$$(x^\lambda, y^i, y^i_\lambda, \dot x^\lambda, \dot y^i, \dot y^i_\lambda) \circ r = (x^\lambda, y^i, y^i_\lambda, \dot x^\lambda, \dot y^i, (\dot y^i)_\lambda - y^i_\mu \dot x^\mu_\lambda),$$

where J^1TY is the jet manifold of the fibred manifold $TY \to X$. In particular, there exists the canonical isomorphism

$$VJ^1Y = J^1VY, \qquad \dot y^i_\lambda = (\dot y^i)_\lambda, \tag{29}$$

where J^1VY is the jet manifold of the fibred manifold $VY \to X$ and VJ^1Y is the vertical tangent bundle of the fibred manifold $J^1Y \to X$. As a consequence, the jet lift (28) of a vertical vector field u on a fibred manifold $Y \to X$ consists with its first order jet prolongation

$$\overline{u} = J^1u = u^i\partial_i + (\partial_\lambda u^i + y^j_\lambda \partial_j u^i)\partial^\lambda_i$$

to a vertical vector field on the fibred jet manifold $J^1Y \to X$.

Given a second order jet manifold J^2Y of Y, we have (i) the fibred morphism

$$r_2 : J^2TY \to TJ^2Y,$$

$$(\dot y^i_\lambda, \dot y^i_{\lambda\alpha}) \circ r_2 = ((\dot y^i)_\lambda - y^i_\mu \dot x^\mu_\lambda, (\dot y^i)_{\lambda\alpha} - y^i_\mu \dot x^\mu_{\lambda\alpha} - y^i_{\lambda\mu}\dot x^\mu_\alpha),$$

and (ii) the canonical isomorphism

$$VJ^2Y = J^2VY$$

where J^2VY is the second order jet manifold of the fibred manifold $VY \to X$ and VJ^2Y is the vertical tangent bundle of the fibred manifold $J^2Y \to X$.

As a consequence, every vector field u on a fibred manifold $Y \to X$ admits the second order jet lift to the projectable vector field

$$\overline{u} = r_2 \circ J^2u : J^2Y \to TJ^2Y.$$

In particular, if

$$u = u^\lambda \partial_\lambda + u^i\partial_i$$

is a projectable vector field on Y, its second order jet lift reads

$$\begin{aligned} \overline{u} = {} & u^\lambda\partial_\lambda + u^i\partial_i + (\partial_\lambda u^i + y^j_\lambda\partial_j u^i - y^i_\mu \partial_\lambda u^\mu)\partial^\lambda_i + \\ & [(\partial_\alpha + y^j_\alpha\partial_j + y^j_{\beta\alpha}\partial^\beta_j)(\partial_\lambda + y^k_\lambda\partial_k)u^i - y^i_\mu \dot x^\mu_{\lambda\alpha} - y^i_{\lambda\mu}\dot x^\mu_\alpha]\partial^{\lambda\alpha}_i. \end{aligned} \tag{30}$$

Given a k-order jet manifold J^kY of Y, we have the fibred morphism

$$r_k : J^kTY \to TJ^kY$$

and the canonical isomorphism

$$VJ^kY = J^kVY$$

where J^kVY is the k-order jet manifold of the fibred manifold $VY \to X$ and VJ^kY is the vertical tangent bundle of the fibred manifold $J^kY \to X$. As a consequence, every vector field u on a fibred manifold $Y \to X$ has the k-order jet lift to the projectable vector field

$$\begin{aligned}
&\overline{u} = r_k \circ J^k u : J^kY \to TJ^kY, \\
&\overline{u} = u^\lambda \partial_\lambda + u^i \partial_i + u^i_\Lambda \partial^\Lambda_i, \\
&u^i_{\Lambda+\lambda} = \widehat{\partial}_\lambda u^i_\Lambda - y^i_{\Lambda+\mu} u^\mu,
\end{aligned} \tag{31}$$

on J^kY where

$$\widehat{\partial}_\lambda = (\partial_\lambda + y^i_{\Sigma+\lambda}\partial^\Sigma_i), \qquad 0 \leq \mid \Sigma \mid \leq k. \tag{32}$$

The expression (31) is the k-order generalization of the expressions (28) and (30).

Algebraic structure of a bundle $Y \to X$ also has the jet prolongation to the jet bundle $J^1Y \to X$ due to the jet prolongations of the corresponding morphisms.

If Y is a vector bundle, $J^1Y \to X$ does as well. Let Y be a vector bundle and $\langle\rangle$ the linear fibred morphism

$$\begin{aligned}
&\langle\rangle : Y \underset{X}{\times} Y^* \underset{X}{\to} X \times \mathbf{R}, \\
&r \circ \langle\rangle = y^i y_i.
\end{aligned}$$

The jet prolongation of $\langle\rangle$ is the linear fibred morphism

$$\begin{aligned}
&J^1\langle\rangle : J^1Y \underset{X}{\times} J^1Y^* \underset{X}{\to} T^*X \times \mathbf{R}, \\
&\dot{x}_\mu \circ J^1\langle\rangle = y^i_\mu y_i + y^i y_{i\mu}.
\end{aligned}$$

Let $Y \to X$ and $Y' \to X$ be vector bundles and $\otimes$ the bilinear fibred morphism

$$\begin{aligned}
&\otimes : Y \underset{X}{\times} Y' \underset{X}{\to} Y \underset{X}{\otimes} Y', \\
&y^{ik} \circ \otimes = y^i y^k.
\end{aligned}$$

The jet prolongation of $\otimes$ is the bilinear fibred morphism

$$\begin{aligned}
&J^1\otimes : J^1Y \underset{X}{\times} J^1Y' \underset{X}{\to} J^1(Y \underset{X}{\otimes} Y'), \\
&y^{ik}_\mu \circ J^1\otimes = y^i_\mu y^k + y^i y^k_\mu.
\end{aligned}$$

If Y is an affine bundle modelled on a vector bundle $\overline{Y}$, then $J^1Y \to X$ is an affine bundle modelled on the vector bundle $J^1\overline{Y} \to X$.

Proposition 1.3: There exist the following bundle monomorphisms:

(i) the contact map

$$\begin{aligned} &\lambda : J^1Y \underset{Y}{\to} T^*X \underset{Y}{\otimes} TY, \qquad (33)\\ &\lambda = dx^\lambda \otimes \widehat{\partial}_\lambda = dx^\lambda \otimes (\partial_\lambda + y^i_\lambda \partial_i), \end{aligned}$$

(ii) the complementary morphism

$$\begin{aligned} &\theta_1 : J^1Y \underset{Y}{\to} T^*Y \underset{Y}{\otimes} VY, \qquad (34)\\ &\theta_1 = \widehat{d}y^i \otimes \partial_i = (dy^i - y^i_\lambda dx^\lambda) \otimes \partial_i. \end{aligned}$$

These monomorphisms able us to handle the jets as familiar tangent-valued forms.

The canonical morphisms (33) and (34) define the bundle monomorphisms

$$\widehat{\lambda} : J^1Y \underset{X}{\times} TX \ni \partial_\lambda \mapsto \widehat{\partial}_\lambda \in J^1Y \underset{Y}{\times} TY, \qquad (35)$$

$$\widehat{\theta}_1 : J^1Y \underset{Y}{\times} V^*Y \ni dy^i \mapsto \widehat{d}y^i \in J^1Y \underset{Y}{\times} T^*Y, \qquad (36)$$

The morphism (35) yields the canonical horizontal splitting of the pullback

$$J^1Y \underset{Y}{\times} TY = \widehat{\lambda}(TX) \underset{J^1Y}{\oplus} VY, \qquad (37)$$

$$\dot{x}^\lambda \partial_\lambda + \dot{y}^i \partial_i = \dot{x}^\lambda(\partial_\lambda + y^i_\lambda \partial_i) + (\dot{y}^i - \dot{x}^\lambda y^i_\lambda)\partial_i.$$

Accordingly, the morphism (36) yields the dual canonical horizontal splitting of the pullback

$$J^1Y \underset{Y}{\times} T^*Y = T^*X \underset{J^1Y}{\oplus} \widehat{\theta}_1(V^*Y), \qquad (38)$$

$$\dot{x}_\lambda dx^\lambda + \dot{y}_i dy^i = (\dot{x}_\lambda + \dot{y}_i y^i_\lambda) dx^\lambda + \dot{y}_i(dy^i - y^i_\lambda dx^\lambda).$$

In other words, over J^1Y, we have the canonical horizontal splittings of the tangent and cotangent bundles TY and T^*Y and the corresponding splittings of the exact sequences (7a) and (7b).

In particular, one gets the canonical horizontal splittings of

(i) a projectable vector field

$$\begin{aligned} u &= u^\lambda \partial_\lambda + u^i \partial_i = u_H + u_V\\ &= u^\lambda(\partial_\lambda + y^i_\lambda \partial_i) + (u^i - u^\lambda y^i_\lambda)\partial_i, \qquad (39) \end{aligned}$$

(ii) an exterior 1-form

$$\begin{aligned} \sigma &= \sigma_\lambda dx^\lambda + \sigma^i dy^i\\ &= (\sigma_\lambda + y^i_\lambda \sigma_i) dx^\lambda + \sigma_i(dy^i - y^i_\lambda dx^\lambda), \end{aligned}$$

(iii) a tangent-valued projectable horizontal form

$$\begin{aligned}\phi &= dx^{\lambda_1}\wedge\ldots\wedge dx^{\lambda_r}\otimes(\phi^\mu_{\lambda_1\ldots\lambda_r}\partial_\mu+\phi^i_{\lambda_1\ldots\lambda_r}\partial_i)\\ &= dx^{\lambda_1}\wedge\ldots\wedge dx^{\lambda_r}\otimes[\phi^\mu_{\lambda_1\ldots\lambda_r}(\partial_\mu+y^i_\mu\partial_i)\\ &\qquad +(\phi^i_{\lambda_1\ldots\lambda_r}-\phi^\mu_{\lambda_1\ldots\lambda_r}y^i_\mu)\partial_i],\end{aligned}$$

(iv) the canonical 1-form

$$\begin{aligned}\theta_Y &= dx^\lambda\otimes\partial_\lambda+dy^i\otimes\partial_i\\ &= \lambda+\theta_1 = dx^\lambda\otimes\widehat{\partial}_\lambda+\widehat{d}y^i\otimes\partial_i\\ &\qquad = dx^\lambda\otimes(\partial_\lambda+y^i_\lambda\partial_i)+(dy^i-y^i_\lambda dx^\lambda)\otimes\partial_i.\end{aligned}\tag{40}$$

As an immediate consequence of the splitting (40), we get the canonical horizontal splitting of the exterior differential

$$d=d_{\theta_Y}=d_H+d_V=d_\lambda+d_{\theta_1}.\tag{41}$$

Its components d_H and d_V act on the pullbacks of horizontal exterior forms

$$\phi_{\lambda_1\ldots\lambda_r}(y)dx^{\lambda_1}\wedge\ldots\wedge dx^{\lambda_r}$$

on a fibred manifold $Y\to X$ by π_{01} onto J^1Y. In this case, d_H makes the sense of the total differential

$$\begin{aligned}&d_H\phi_{\lambda_1\ldots\lambda_r}(y)dx^{\lambda_1}\wedge\ldots\wedge dx^{\lambda_r}\\ &\qquad=(\partial_\mu+y^i_\mu\partial_i)\phi_{\lambda_1\ldots\lambda_r}(y)dx^\mu\wedge dx^{\lambda_1}\wedge\ldots\wedge dx^{\lambda_r},\end{aligned}$$

whereas d_V is the vertical differential

$$\begin{aligned}&d_V\phi_{\lambda_1\ldots\lambda_r}(y)dx^{\lambda_1}\wedge\ldots\wedge dx^{\lambda_r}\\ &\qquad=\partial_i\phi_{\lambda_1\ldots\lambda_r}(y)(dy^i-y^i_\mu dx^\mu)\wedge dx^{\lambda_1}\wedge\ldots\wedge dx^{\lambda_r}.\end{aligned}$$

If $\phi=\widetilde{\phi}\omega$ is an exterior horizontal density on $Y\to X$, we have

$$d\phi=d_V\phi=\partial_i\widetilde{\phi}dy^i\wedge\omega.$$

There exist the following second order generalizations of the contact map (33) and the complementary morphism (34) to the second order jet manifold J^2Y:

$$\begin{aligned}(i)\quad &\lambda: J^2Y\underset{J^1Y}{\to}T^*X\underset{J^1Y}{\otimes}TJ^1Y,\\ &\lambda=dx^\lambda\otimes\widehat{\partial}_\lambda=dx^\lambda\otimes(\partial_\lambda+y^i_\lambda\partial_i+y^i_{\mu\lambda}\partial^\mu_i),\end{aligned}\tag{42}$$

$$\begin{aligned}(ii)\quad &\theta_1: J^2Y\underset{J^1Y}{\to}T^*J^1Y\underset{J^1Y}{\otimes}VJ^1Y,\\ &\theta_1=(dy^i-y^i_\lambda dx^\lambda)\otimes\partial_i+(dy^i_\mu-y^i_{\mu\lambda}dx^\lambda)\otimes\partial^\mu_i.\end{aligned}\tag{43}$$

The contact map (42) defines the canonical horizontal splitting of the exact sequence

$$0 \to VJ^1Y \underset{J^1Y}{\hookrightarrow} TJ^1Y \underset{J^1Y}{\to} J^1Y \underset{X}{\times} TX \to 0.$$

In particular, we get the canonical horizontal splitting of a projectable vector field $\overline{u}$ on J^1Y over J^2Y:

$$\begin{aligned} \overline{u} = u_H + u_V = u^\lambda[\partial_\lambda + y^i_\lambda + y^i_{\mu\lambda}] \\ +[(u^i - y^i_\lambda u^\lambda)\partial_i + (u^i_\mu - y^i_{\mu\lambda}u^\lambda)\partial^\mu_i]. \end{aligned} \tag{44}$$

Using the morphisms (42) and (43), we can still obtain the horizontal splittings of the canonical tangent-valued 1-form θ_{J^1Y} on J^1Y and, as a result, the horizontal splitting of the exterior differential which are similar the horizontal splittings (40) and (41):

$$\begin{aligned} &\theta_{J^1Y} = dx^\lambda \otimes \partial_\lambda + dy^i \otimes \partial_i + dy^i_\mu \otimes \partial^\mu_i = \lambda + \theta_1, \\ &d = d_{\theta_{J^1Y}} = d_\lambda + d_{\theta_1} = d_H + d_V. \end{aligned} \tag{45}$$

The contact maps (33) and (42) are the particular cases of the monomorphism

$$\begin{aligned} &\lambda : J^{k+1}Y \to T^*X \underset{J^kY}{\otimes} TJ^kY, \\ &\lambda = dx^\lambda \otimes (\partial_\lambda + y^i_{\Lambda+\lambda}\partial^\Lambda_i), \qquad 0 \leq\mid \Lambda \mid\leq k. \end{aligned} \tag{46}$$

The k-order contact map (46) sets up the canonical horizontal splitting of the exact sequence

$$0 \to VJ^kY \hookrightarrow TJ^kY \to J^kY \underset{X}{\times} TX \to 0.$$

In particular, we get the canonical horizontal splitting of a projectable vector field $\overline{u}$ on $J^kY \to X$ over $J^{k+1}Y$:

$$\overline{u} = u_H + u_V = u^\lambda(\partial_\lambda + y^i_{\Lambda+\lambda}\partial^\Lambda_i) + (u^i_\Lambda - y^i_{\Lambda+\lambda})\partial^\Lambda_i, \qquad 0 \leq\mid \Lambda \mid\leq k.$$

This splitting is the k-order generalization of the splittings (39) and (30).

Definition 1.4: The repeated jet manifold J^1J^1Y is defined to be the first order jet manifold of the fibred jet manifold $J^1Y \to X$.

Given the coordinates (24) of J^1Y, the repeated jet manifold J^1J^1Y is provided with the adapted coordinates

$$(x^\lambda, y^i, y^i_\lambda, y^i_{(\mu)}, y^i_{\lambda\mu}). \tag{47}$$

There are the following two bundles:

$$\begin{aligned} (i) \quad & \pi_{11} : J^1J^1Y \to J^1Y, && (48) \\ & y^i_\lambda \circ \pi_{11} = y^i_\lambda, \\ (ii) \quad & J^1\pi_{01} : J^1J^1Y \to J^1Y, && (49) \\ & y^i_\lambda \circ J^1\pi_{01} = y^i_{(\lambda)}. \end{aligned}$$

Their affine difference over Y yields the Spencer bundle morphism

$$\delta = J^1\pi_{01} - \pi_{11} : J^1J^1Y \underset{Y}{\to} T^*X \underset{Y}{\otimes} VY,$$

$$(x^\lambda, y^i, \dot{x}_\lambda \otimes \dot{y}^i) \circ \delta = (x^\lambda, y^i, y^i_{(\lambda)} - y^i_\lambda).$$

The kernel of this morphism is the sesquiholonomic affine subbundle

$$\widehat{J}^2Y \to J^1Y \tag{50}$$

of the bundles (48) and (49). This subbundle is characterized by the coordinate condition $y^i_{(\lambda)} = y^i_\lambda$. It is modelled on the vector bundle

$$\overset{2}{\otimes} T^*X \underset{J^1Y}{\otimes} VY.$$

Given the coordinates (47) of J^1J^1Y, the sesquiholonomic jet manifold $\widehat{J}^2Y$ is provided with the adapted coordinates $(x^\lambda, y^i, y^i_\lambda, y^i_{\lambda\mu})$. The second order jet manifold J^2Y is the affine subbundle of the bundle (50) given by the coordinate condition $y^i_{\lambda\mu} = y^i_{\mu\lambda}$. It is modelled on the vector bundle

$$\overset{2}{\vee} T^*X \underset{J^1Y}{\otimes} VY.$$

We have the following affine bundle monomorphisms

$$J^2Y \hookrightarrow \widehat{J}^2Y \hookrightarrow J^1J^1Y$$

over J^1Y and the canonical splitting

$$\widehat{J}^2Y = J^2Y \underset{J^1Y}{\oplus} (\overset{2}{\wedge} T^*X \underset{Y}{\otimes} VY), \tag{51}$$

$$y^i_{\lambda\mu} = \frac{1}{2}(y^i_{\lambda\mu} + y^i_{\mu\lambda}) + (\frac{1}{2}(y^i_{\lambda\mu} - y^i_{\mu\lambda}).$$

Let Φ be a fibred morphism of a fibred manifold $Y \to X$ to a fibred manifold $Y' \to X$ over a diffeomorphism of X. Let $J^1\Phi$ be its first order jet prolongation (26). One can consider the first order jet prolongation $J^1J^1\Phi$ of the fibred morphism $J^1\Phi$. The restriction of the morphism $J^1J^1\Phi$ to the second order jet manifold J^2Y of Y consists with the second order jet prolongation $J^2\Phi$ of a fibred morphism Φ.

In particular, the repeated jet prolongation J^1J^1s of a section s of $Y \to X$ is a section of the fibred manifold $J^1J^1Y \to X$. It takes its values into J^2Y and consists with the second order jet prolongation J^2s of s:

$$(J^1J^1s)(x) = (J^2s)(x) = j^2_x s.$$

Given the fibred jet manifold $J^kY \to X$, let us consider the repeated jet manifold J^1J^kY provided with the adapted coordinates $(x^\mu, y^i_\Lambda, y^i_{\Lambda\lambda})$. Just as in the case of $k = 2$, there exist two fibred morphisms of J^1J^kY to $J^1J^{k-1}Y$ over X. Their difference over $J^{k-1}Y$ is the k-order Spencer morphism

$$J^1J^kY \to T^*X \underset{J^{k-1}Y}{\otimes} VJ^{k-1}Y$$

where $VJ^{k-1}Y$ is the vertical tangent bundle of the fibred manifold $J^{k-1}Y \to X$. Its kernel is the sesquiholonomic subbundle $\widehat{J}^{k+1}Y$ of the bundle $J^{k+1}Y \to Y$. It is endowed with the coordinates $(x^\mu, y^i_\Lambda, y^i_{\Lambda\mu})$, $\mid \Lambda \mid= k$.

Proposition 1.5: There exist the fibred monomorphisms

$$J^kY \hookrightarrow \widehat{J}^kY \hookrightarrow J^1J^{k-1}Y \tag{52}$$

and the canonical splitting

$$\widehat{J}^{k+1}Y = J^{k+1}Y \underset{J^kY}{\oplus} (T^*X \wedge \overset{k-1}{\vee} T^*X \underset{Y}{\otimes} VY), \tag{53}$$

$$(x^\mu, y^i_\Lambda, y^i_{\Lambda\mu}) = (x^\mu, y^i_\Lambda, y^i_{(\Lambda\mu)} + y^i_{[\Lambda\mu]}).$$

We have the following integrability condition.

Lemma 1.6: Let $\overline{s}$ be a section of the fibred manifold $J^kY \to X$. Then, the following conditions are equivalent:

(i) $\overline{s} = J^ks$ where s is a section of $Y \to X$,
(ii) $J^1\overline{s} : X \to \widehat{J}^{k+1}Y$,
(iii) $J^1\overline{s} : X \to J^{k+1}Y$.

Building on Proposition 1.5 and Lemma 1.6, we now describe reduction of higher order differential operators to the first order ones.

Let Y and Y' be fibred manifolds over X and J^kY the k-order jet manifold of Y.

Definition 1.7: A fibred morphism

$$\mathcal{E} : J^kY \underset{X}{\to} Y' \tag{54}$$

is called the k-order differential operator (of class C^∞) on Y. It sends every section s of the fibred manifold Y to the section $\mathcal{E} \circ J^ks$ of the fibred manifold Y'.

Proposition 1.8: Given a fibred manifold Y, every first order differential operator

$$\mathcal{E}'' : J^1J^{k-1}Y \underset{X}{\to} Y' \tag{55}$$

on $J^{k-1}Y$ defines the k-order differential operator $\mathcal{E} = \mathcal{E}'' \mid_{J^kY}$ on Y. Conversely, if a first order differential operator on $J^{k-1}Y$ exists, any k-rder differential operator (54) on Y can

be represented by the restriction $\mathcal{E}''\mid_{J^kY}$ of some first order differential operator (55) on $J^{k-1}Y$ to the k-order jet manifold J^kY.

Proof. Because of the monomorphism (52) every fibred morphism $J^kY \to Y'$ can be extended to a fibred morphism $J^1J^{k-1}Y \to Y'$.

In particular, every k-order differential operator (54) yields the morphism

$$\mathcal{E}' = \mathcal{E} \circ \mathrm{pr}_2 : \widehat{J}^kY \underset{X}{\to} Y' \tag{56}$$

where $\mathrm{pr}_2 : \widehat{J}^kY \to J^kY$ is the surjection corresponding to the canonical splitting (53). It follows that, for every section s of a fibred manifold $Y \to X$, we have the equality

$$\mathcal{E}' \circ J^1J^{k-1}s = \mathcal{E} \circ J^ks.$$

We call $\mathcal{E}'$ (56) the sesquiholonomic differential operator and consider extensions of a k-order differential operator $\mathcal{E}$ (54) to first order differential operators (55) only through its extension to the sequiholonomic differential operator (56).

Let $\overline{s}$ be a section of the fibred $(k-1)$-order jet manifold $J^{k-1}Y \to X$ such that its first order jet prolongation $J^1\overline{s}$ takes its values into the sesquiholonomic jet manifold $\widehat{J}^kY$. In virtue of Lemma 1.6, there exists a section s of $Y \to X$ such that $\overline{s} = J^{k-1}s$ and

$$\mathcal{E}' \circ J^1\overline{s} = \mathcal{E} \circ J^ks. \tag{57}$$

Let $Y' \to X$ be a composite manifold $Y' \to Y \to X$ where $Y' \to Y$ is a vector bundle. Let the k-order differential operator (54) on a fibred manifold Y be a fibred morphism over Y. By $\mathrm{Ker}\,\mathcal{E}$ is meant the preimage $\mathcal{E}^{-1}(\widehat{0}(Y))$ where $\widehat{0}$ is the canonical zero section of $Y' \to Y$. We say that a section s of Y satisfies the corresponding system of k-order differential equations if

$$J^ks(X) \subset \mathrm{Ker}\,\mathcal{E}. \tag{58}$$

Let a k-order differential operator $\mathcal{E}$ on $Y \to X$ be extended to a first order differential operator $\mathcal{E}''$ on $J^{k-1}Y$. Let $\overline{s}$ be a section of the fibred manifold $J^{k-1}Y \to X$. We shall say that $\overline{s}$ is a sesquiholonomic solution of the corresponding system of first order differential equations if

$$J^1\overline{s}(X) \subset \mathrm{Ker}\,\mathcal{E}'' \cap \widehat{J}^kY. \tag{59}$$

Proposition 1.9: The system of the k-order differential equations (58) and the system of the first order differential equations (59) are equivalent to each other.

Proof. In virtue of the relation (57), every solution s of the equations (58) yields the solution

$$\overline{s} = J^{k-1}s \tag{60}$$

of the equations (59). Every solution of the equations (59) takes the form (60) where s is a solution of the equations (58).

3 General Connections

The most of existent differential geometric methods in field theory is based on principal bundles and principal connections. We follow the general notion of connections as sections of jet bundles $J^1Y \to Y$ without appealing to transformation groups.

Given a fibred manifold $Y \to X$, the canonical horizontal splittings (37) and (38) of the tangent and cotangent bundles TY and T^*Y of Y over the jet bundle $J^1Y \to Y$ enable us to get the splittings of the exact sequences (7a) and (7b) by means of a section of this jet bundle.

Definition 1.10: A first order jet field (or simply a jet field) on a fibred manifold $Y \to X$ is defined to be a section Γ of the affine jet bundle $J^1Y \to Y$. A first order connection Γ on a fibred manifold Y is defined to be a global jet field

$$\begin{aligned} &\Gamma : Y \to J^1Y, \\ &(x^\lambda, y^i, y^i_\lambda) \circ \Gamma = (x^\lambda, y^i, \Gamma^i_\lambda(y)). \end{aligned} \tag{61}$$

By means of the contact map λ (33), every connection Γ (61) on a fibred manifold $Y \to X$ can be represented by a projectable tangent-valued horizontal 1-form $\lambda \circ \Gamma$ on Y which we denote by the same symbol

$$\begin{aligned} &\Gamma = dx^\lambda \otimes (\partial_\lambda + \Gamma^i_\lambda(y)\partial_i), \qquad (62) \\ &\Gamma'^i_\lambda = (\frac{\partial y'^i}{\partial y^j}\Gamma^j_\mu + \frac{\partial y'^i}{\partial x^\mu})\frac{\partial x^\mu}{\partial x'^\lambda}. \end{aligned}$$

Substituting a connection Γ (62) into the canonical horizontal splittings (37) and (38), we obtain the familiar horizontal splittings

$$\begin{aligned} &\dot x^\lambda \partial_\lambda + \dot y^i \partial_i = \dot x^\lambda(\partial_\lambda + \Gamma^i_\lambda \partial_i) + (\dot y^i - \dot x^\lambda \Gamma^i_\lambda)\partial_i, \\ &\dot x_\lambda dx^\lambda + \dot y_i dy^i = (\dot x_\lambda + \Gamma^i_\lambda \dot y_i)dx^\lambda + \dot y_i(dy^i - \Gamma^i_\lambda dx^\lambda) \end{aligned} \tag{63}$$

of the tangent and cotangent bundles TY and T^*Y with respect to a connection Γ on Y. Conversely, every horizontal splitting (63) determines a tangent-valued form (62) and, consequently, a global jet field on $Y \to X$.

Since the affine jet bundle $J^1Y \to Y$ is modelled on the vector bundle (25), connections on a fibred manifold Y constitute an affine space modelled on the linear space of soldering forms on Y. It follows that, if Γ is a connection and

$$\sigma = \sigma^i_\lambda dx^\lambda \otimes \partial_i$$

is a soldering form on a fibred manifold Y, then

$$\Gamma + \sigma = dx^\lambda \otimes [\partial_\lambda + (\Gamma^i_\lambda + \sigma^i_\lambda)\partial_i]$$

is a connection on Y. Conversely, if Γ and Γ' are connections on a fibred manifold Y, then

$$\Gamma - \Gamma' = (\Gamma^i_\lambda - \Gamma'^i_\lambda) dx^\lambda \otimes \partial_i$$

is a soldering form on Y.

For instance, let $Y \to X$ be a vector bundle. A linear connection on Y reads

$$\Gamma = dx^\lambda \otimes [\partial_\lambda + \Gamma^i{}_{j\lambda}(x) y^j \partial_i]. \tag{64}$$

One introduces the following basic forms involving a connection Γ and a soldering form σ on a fibred manifold Y:

(i) the curvature of Γ:

$$\begin{aligned} R = \frac{1}{2} d_\Gamma \Gamma = \frac{1}{2} R^i_{\lambda\mu} dx^\lambda \wedge dx^\mu \otimes \partial_i = \\ \frac{1}{2}(\partial_\lambda \Gamma^i_\mu - \partial_\mu \Gamma^i_\lambda + \Gamma^j_\lambda \partial_j \Gamma^i_\mu - \Gamma^j_\mu \partial_j \Gamma^i_\lambda) dx^\lambda \wedge dx^\mu \otimes \partial_i; \end{aligned} \tag{65}$$

(ii) the torsion of Γ with respect to σ:

$$\begin{aligned} \Omega = d_\sigma \Gamma = d_\Gamma \sigma = \frac{1}{2} \Omega^i_{\lambda\mu} dx^\lambda \wedge dx^\mu \otimes \partial_i = \\ (\partial_\lambda \sigma^i_\mu + \Gamma^j_\lambda \partial_j \sigma^i_\mu - \partial_j \Gamma^i_\lambda \sigma^j_\mu) dx^\lambda \wedge dx^\mu \otimes \partial_i; \end{aligned} \tag{66}$$

(ii) the soldering curvature of σ:

$$\begin{aligned} \varepsilon = \frac{1}{2} d_\sigma \sigma = \frac{1}{2} \varepsilon^i_{\lambda\mu} dx^\lambda \wedge dx^\mu \otimes \partial_i = \\ \frac{1}{2}(\sigma^j_\lambda \partial_j \sigma^i_\mu - \sigma^j_\mu \partial_j \sigma^i_l) dx^\lambda \wedge dx^\mu \otimes \partial_i. \end{aligned} \tag{67}$$

In particular, the curvature (65) of the linear connection (64) reads

$$\begin{aligned} & R^i_{\lambda\mu}(y) = R^i{}_{j\lambda\mu}(x) y^j, \\ & R^i{}_{j\lambda\mu} = \partial_\lambda \Gamma^i{}_{j\mu} - \partial_\mu \Gamma^i{}_{j\lambda} + \Gamma^k{}_{j\lambda} \Gamma^i{}_{k\mu} - \Gamma^k{}_{j\mu} \Gamma^i{}_{k\lambda}. \end{aligned} \tag{68}$$

A connection Γ on a fibred manifold $Y \to X$ yields the affine bundle morphism

$$\begin{aligned} & D_\Gamma : J^1 Y \ni z \mapsto z - \Gamma(\pi_{01}(z)) \in T^* X \underset{Y}{\otimes} VY, \\ & D_\Gamma = (y^i_\lambda - \Gamma^i_\lambda) dx^\lambda \otimes \partial_i. \end{aligned} \tag{69}$$

It is called the covariant differential. The corresponding covariant derivative of sections s of Y reads

$$\nabla_\Gamma s = D_\Gamma \circ J^1 s = (\partial_\lambda s^i - (\Gamma \circ s)^i_\lambda) dx^\lambda \otimes \partial_i \tag{70}$$

A section s of a fibred manifold Y is called an integral section for a connection Γ on Y if

$$\Gamma\circ s = J^1 s, \tag{71}$$

that is, $\nabla_\Gamma s = 0$.

Now, we consider some particular properties of linear connections on vector bundles.

Let $Y\to X$ be a vector bundle and Γ a linear connection on Y. On the dual vector bundle $Y^*\to X$, there exists the dual connection Γ^* is called the dual connection to Γ given by the coordinate expression

$$\Gamma^*_{i\lambda} = -\Gamma^j{}_{i\lambda}(x)y_j.$$

For instance, a linear connection K on the tangent bundle TX of a manifold X and the dual connection K^* to K on the cotangent bundle T^*X read

$$\begin{aligned} &K^\alpha_\lambda = K^\alpha{}_{\nu\lambda}(x)\dot x^\nu,\\ &K^*_{\alpha\lambda} = -K^\nu{}_{\alpha\lambda}(x)\dot x_\nu, \end{aligned} \tag{72}$$

Let Y and Y' be vector bundles over X. Given linear connections Γ and Γ' on Y and Y' respectively, the tensor product connection $\Gamma\otimes\Gamma'$ on the tensor product

$$Y\underset{X}{\otimes} Y'\to X$$

is defined. It takes the coordinate form

$$(\Gamma\otimes\Gamma')^{ik}_\lambda = \Gamma^i{}_{j\lambda}y^{jk} + \Gamma'^k{}_{j\lambda}y^{ij}.$$

The construction of the dual connection and the tensor product connection can be extended to connections on composite manifolds (4) when $Y\to\Sigma$ is a vector bundle.

Let $Y\to\Sigma\to X$ be the composite manifold (4). Let $J^1\Sigma$, J^1Y_Σ and J^1Y the first order jet manifolds of $\Sigma\to X$, $Y\to\Sigma$ and $Y\to X$ respectively. Given fibred coordinates (x^λ,σ^m,y^i) (5) of Y, the corresponding adapted coordinates of the jet manifolds $J^1\Sigma$, J^1Y_Σ and J^1Y are

$$\begin{aligned} &(x^\lambda,\sigma^m,\sigma^m_\lambda),\\ &(x^\lambda,\sigma^m,y^i,\widetilde y^i_\lambda,y^i_m),\\ &(x^\lambda,\sigma^m,y^i,\sigma^m_\lambda,y^i_\lambda). \end{aligned} \tag{73}$$

We say that a connection

$$A = dx^\lambda\otimes[\partial_\lambda + \Gamma^m_\lambda(\sigma)\partial_m + A^i_\lambda(y)\partial_i], \qquad \sigma = \pi_{Y\Sigma}(y),$$

on a composite manifold $Y\to\Sigma\to X$ is projectable to a connection

$$\Gamma = dx^\lambda\otimes[\partial_\lambda + \Gamma^m_\lambda(\sigma)\partial_m]$$

on the fibred manifold $\Sigma \to X$, if there exists the commutative diagram

$$\begin{tikzcd}
J^1Y \arrow[r, "J^1\pi_{Y\Sigma}"] & J^1\Sigma \\
Y \arrow[u, "A"] \arrow[r, "\pi_{Y\Sigma}"'] & \Sigma \arrow[u, "\Gamma"']
\end{tikzcd}$$

Let $Y \to \Sigma$ be a vector bundle and

$$A = dx^\lambda \otimes [\partial_\lambda + \Gamma^m_\lambda(\sigma)\partial_m + A^i{}_{j\lambda}(\sigma)y^j\partial_i]$$

a linear morphism over Γ. Let $Y^* \to \Sigma \to X$ be a composite manifold where $Y^* \to \Sigma$ is the vector bundle dual to $Y \to \Sigma$. On $Y^* \to X$, there exists the dual connection

$$A^* = dx^\lambda \otimes [\partial_\lambda + \Gamma^m_\lambda(\sigma)\partial_m - A^j{}_{i\lambda}(\sigma)y_j\partial^i] \tag{74}$$

projectable to Γ.

Let $Y \to \Sigma \to X$ and $Y' \to \Sigma \to X$ be composite manifolds where $Y \to \Sigma$ and $Y' \to \Sigma$ are vector bundles. Let A and A' be connections on Y and Y' respectively which are projectable to the same connection Γ on the fibred manifold $\Sigma \to X$. On the tensor product

$$Y \underset{\Sigma}{\otimes} Y' \to X,$$

there exists the tensor product connection

$$A \otimes A' = dx^\lambda \otimes (\partial_\lambda + \Gamma^m_\lambda\partial_m + (A^i{}_{j\lambda}y^{jk} + A'^k{}_{j\lambda}y^{ij})\partial_{ik}) \tag{75}$$

projectable to Γ.

In particular, let $Y \to X$ be a fibred manifold and Γ a connection on Y. The vertical tangent morphism $V\Gamma$ to Γ defines the connection

$$\begin{aligned}
&V\Gamma : VY \to VJ^1Y = J^1VY, \\
&V\Gamma = dx^\lambda \otimes (\partial_\lambda + \Gamma^i_\lambda \frac{\partial}{\partial y^i} + \partial_j\Gamma^i_\lambda \dot y^j \frac{\partial}{\partial \dot y^i}),
\end{aligned} \tag{76}$$

on the composite manifold $VY \to Y \to X$ due to the canonical isomorphism (29). The connection $V\Gamma$ is projectable to the connection Γ on Y, and it is a linear bundle morphism over Γ:

$$\begin{tikzcd}
VY \arrow[r, "V\Gamma"] \arrow[d] & J^1VY \arrow[d] \\
Y \arrow[r, "\Gamma"] & J^1Y
\end{tikzcd}$$

The connection (76) yields the connection

$$V^*\Gamma = dx^\lambda \otimes (\partial_\lambda + \Gamma^i_\lambda \frac{\partial}{\partial y^i} - \partial_j\Gamma^i_\lambda \dot y_i \frac{\partial}{\partial \dot y_j}) \tag{77}$$

on the composite manifold $V^*Y \to Y \to X$ which is the dual connection to $V\Gamma$ over Γ.

Now, we consider second-order connections.

Definition 1.11: A second order jet field (resp. a second order connection) $\overline{\Gamma}$ on a fibred manifold $Y \to X$ is defined to be a first order jet field (resp. a first order connection) on the fibred jet manifold $J^1Y \to X$, i.e. this is a section (resp. a global section) of the bundle (48).

In the coordinates (47) of the repeated jet manifold J^1J^1Y, a second order jet field $\overline{\Gamma}$ is given by the expression

$$(y^i_\lambda, y^i_{(\mu)}, y^i_{\lambda\mu}) \circ \overline{\Gamma} = (y^i_\lambda, \overline{\Gamma}^i_{(\mu)}, \overline{\Gamma}^i_{\lambda\mu}).$$

Using the contact map (42), one can represent it by the horizontal 1-form

$$\overline{\Gamma} = dx^\mu \otimes (\partial_\mu + \overline{\Gamma}^i_{(\mu)}\partial_i + \overline{\Gamma}^i_{\lambda\mu}\partial^\lambda_i) \tag{78}$$

on the fibred jet manifold $J^1Y \to X$.

A second order jet field $\overline{\Gamma}$ on Y is termed a sesquiholonomic (resp. holonomic) second order jet field if it takes its values into the subbundle $\widehat{J}^2Y$ (resp. J^2Y) of J^1J^1Y. We have the coordinate equality $\overline{\Gamma}^i_{(\mu)} = y^i_\mu$ for a sesquiholonomic second order jet field and the additional equality $\overline{\Gamma}^i_{\lambda\mu} = \overline{\Gamma}^i_{\mu\lambda}$ for a holonomic second order jet field.

Given a first order connection Γ on a fibred manifold $Y \to X$, one can construct a second order connection on Y, that is, a connection on the fibred jet manifold $J^1Y \to X$ as follows.

The first order jet prolongation $J^1\Gamma$ of the connection Γ on Y is a section of the bundle (49), but not the bundle π_{11} (48). Let K^* be a linear symmetric connection (72) on the cotangent bundle T^*X of X:

$$K_{\lambda\mu} = -K^\alpha{}_{\lambda\mu}\dot{x}_\alpha, \qquad K_{\lambda\mu} = K_{\mu\lambda}.$$

There exists the affine fibred morphism

$$r_K : J^1J^1Y \to J^1J^1Y, \qquad r_K \circ r_K = \mathrm{Id}\,_{J^1J^1Y},$$

$$(y^i_\lambda, y^i_{(\mu)}, y^i_{\lambda\mu}) \circ r_K = (y^i_{(\lambda)}, y^i_\mu, y^i_{\mu\lambda} + K^\alpha{}_{\lambda\mu}(y^i_\alpha - y^i_{(\alpha)})).$$

One can verify the following transformation relations of the coordinates (47):

$$y'^i_\mu \circ r_k = y'^i_{(\mu)}, \qquad y'^i_{(\mu)} \circ r_k = y'^i_\mu,$$
$$y'^i_{\lambda\mu} \circ r_k = y'^i_{\mu\lambda} + K'^\alpha{}_{\lambda\mu}(y'^i_\alpha - y'^i_{(\alpha)}).$$

Hence, given a first order connection Γ on a fibred manifold $Y \to X$, we have the second order connection

$$J\Gamma = r_K \circ J^1\Gamma,$$

$$J\Gamma = dx^\mu \otimes [\partial_\mu + \Gamma^i_\mu\partial_i + (\partial_\lambda\Gamma^i_\mu + \partial_j\Gamma^i_\mu y^j_\lambda - K^\alpha{}_{\lambda\mu}(y^i_\alpha - \Gamma^i_\alpha))\partial^\lambda_i], \tag{79}$$

on Y. This is an affine morphism

$$\begin{tikzcd}
J^1Y \arrow[r, "J\Gamma"] \arrow[d, "\pi_{01}"'] & J^1J^1Y \arrow[d, "\pi_{11}"] \\
Y \arrow[r, "\Gamma"'] & J^1Y
\end{tikzcd}$$

over the first order connection Γ.

Note that the curvature R (65) of a first order connection Γ on a fibred manifold $Y \to X$ induces the soldering form

$$\overline{\sigma}_R = R^i_{\lambda\mu} dx^\mu \otimes \partial_i^\lambda \tag{80}$$

on the fibred jet manifold $J^1Y \to X$. Also the torsion (66) of a first order connection Γ with respect to a soldering form σ on $Y \to X$ and the soldering curvature (67) of σ define soldering forms on $J^1Y \to X$.

References

[1] J. Cariñena, M. Crampin and L. Ibort, *Differential Geometry and its Application* **1** (1991) 345.

[2] G. Giachetta and L. Mangiarotti, *International Journal of Theoretical Physics* **29** (1990), 789.

[3] M. Gotay, in *Mechanics, Analysis and Geometry: 200 Years after Lagrange*, ed. M.Francaviglia (Elseiver Science Publishers B.V., 1991) p. 203.

[4] C. Günther, *Journal of Differential Geometry* **25** (1987) 23.

[5] I. Kolář, *Journal of Geometry and Physics* **1** (1984) 127.

[6] B. Kupershmidt, in Lect. Notes in Math., Vol. 775 (Springer, Berlin - New York, 1980) p.162.

[7] L. Mangiarotti and M. Modugno, in *Geometry and Physics*, ed. M. Modugno (Pitagora Editrice, Bologna, 1982) p. 135.

[8] G. Sardanashvily and O. Zakharov, *Differential Geometry and its Applications* **3** (1993) 245.

[9] G. Sardanashvily, *Gauge Theory in Jet Manifolds* (Hadronic Press Inc., Palm Harbor, 1993).

[10] G.Sardanashvily, Multimomentum Hamiltonian Formalism in Field Theory, LaTeX preprint: hep-th/9403172

[11] D. Saunders, *The Geometry of Jet Bundles* (Cambridge Univ. Press, Cambridge, 1989).